



Dietary factors during pregnancy and atopic outcomes in childhood: A systematic review from the European Academy of Allergy and Clinical Immunology

Carina Venter^{1,2} | Carlo Agostoni³ | S. Hasan Arshad^{4,5} |
 Miriam Ben-Abdallah² | George Du Toit^{6,7} | David M. Fleischer^{1,2} |
 Matthew Greenhawt^{1,2} | Deborah H. Glueck⁸ | Marion Groetch⁹ |
 Nonhlanhla Lunjani^{10,11} | Kate Maslin¹² | Alexander Maiorella¹³ | Rosan Meyer¹⁴ |
 Muraro Antonella¹⁵ | Merryn J. Netting^{16,17} | Bright Ibeabughichi Nwaru¹⁸ |
 Debra J. Palmer¹⁹ | Micheala P. Palumbo²⁰ | Graham Roberts^{5,6,21,22} |
 Caroline Roduit^{23,24} | Pete Smith²⁵ | Eva Untersmayr²⁶ |
 Lauren A. Vanderlinden²⁷ | Liam O'Mahony²⁸

¹Section of Allergy and Immunology, University of Colorado School of Medicine, Denver, CO, USA

²Children's Hospital Colorado, Aurora, CO, USA

³Pediatria Media Intensità di Cura Fondazione IRCCS Ca' Granda - Ospedale Maggiore Policlinic, Milan, Italy

⁴Clinical & Experimental Sciences, Faculty of Medicine, University of Southampton, Southampton, UK

⁵The David Hide Asthma and Allergy Centre, Isle of Wight, UK

⁶Department of Paediatric Allergy, Division of Asthma, Allergy and Lung Biology, King's College London, London, UK

⁷Evelina London, Guy's & St Thomas' Hospital, London, UK

⁸Department of Pediatrics, University of Colorado School of Medicine, Denver, CO, USA

⁹Icahn School of Medicine at Mount Sinai, New York, NY, USA

¹⁰University of Zurich, Davos, Switzerland

¹¹University of Cape Town, Cape Town, South Africa

¹²University of Plymouth, Plymouth, UK

¹³University of Chicago, Chicago, IL, USA

¹⁴Imperial College, London, UK

¹⁵Centro di Specializzazione Regionale per lo Studio e la Cura delle Allergie e delle Intolleranze Alimentari presso l'Azienda Ospedaliera, Università di Padova, Padova, Italy

¹⁶Women and Kids Theme, South Australian Health and Medical Research Institute, Adelaide, SA, Australia

¹⁷Discipline of Pediatrics, University of Adelaide, Adelaide, SA, Australia

¹⁸Krefting Research Centre, Institute of Medicine, University of Gothenburg, Gothenburg, Sweden

¹⁹Telethon Kids Institute, University of Western Australia, Perth, WA, Australia

²⁰Lifecourse Epidemiology of Adiposity and Diabetes (LEAD) Center, Colorado School of Public Health, Colorado, USA

²¹NIHR Biomedical Research Centre, University Hospital Southampton NHS Foundation Trust, Southampton, UK

²²Faculty of Medicine, Human Development in Health Academic Units, University of Southampton, Southampton, UK

²³University Children's Hospital Zurich, Switzerland

²⁴Christine Kühne-Center for Allergy Research and Education, Davos, Switzerland

²⁵School of Medicine, Griffith University, Southport, Australia

Abbreviations: AD, Atopic dermatitis; AI, Adequate intake; FA, Food allergy; PC, Prospective cohort study; RDI, Recommended dietary intake.

[Correction added on 18 August 2020, after first online publication: URL for peer review history has been corrected.]

The peer review history for this article is available at <https://publons.com/publon/10.1111/PAI.13303>

© 2020 EAACI and John Wiley and Sons A/S. Published by John Wiley and Sons Ltd.

²⁶Institute of Pathophysiology and Allergy Research, Center for Pathophysiology, Infectiology and Immunology, Medical University of Vienna, Vienna, Austria

²⁷Department of Biostatistics and Informatics, Colorado School of Public Health, Aurora, CO, USA

²⁸Departments of Medicine and Microbiology, APC Microbiome Ireland, National University of Ireland, Cork, Ireland

Correspondence

Carina Venter, Section of Allergy and Immunology, University of Colorado School of Medicine, Denver, CO.

Email: Carina.Venter@childrenscolorado.org

Editor: Rachel Peters

Abstract

Rationale: Allergic diseases are an increasing public health concern, and early life environment is critical to immune development. Maternal diet during pregnancy has been linked to offspring allergy risk. In turn, maternal diet is a potentially modifiable factor, which could be targeted as an allergy prevention strategy. In this systematic review, we focused on non-allergen-specific modifying factors of the maternal diet in pregnancy on allergy outcomes in their offspring.

Methods: We undertook a systematic review of studies investigating the association between maternal diet during pregnancy and allergic outcomes (asthma/wheeze, hay fever/allergic rhinitis/seasonal allergies, eczema/atopic dermatitis (AD), food allergies, and allergic sensitization) in offspring. Studies evaluating the effect of food allergen intake were excluded. We searched three bibliographic databases (MEDLINE, EMBASE, and Web of Science) through February 26, 2019. Evidence was critically appraised using modified versions of the Cochrane Collaboration Risk of Bias tool for intervention trials and the National Institute for Clinical Excellence methodological checklist for cohort and case-control studies and meta-analysis performed from RCTs.

Results: We identified 95 papers: 17 RCTs and 78 observational (case-control, cross-sectional, and cohort) studies. Observational studies varied in design and dietary intakes and often had contradictory findings. Based on our meta-analysis, RCTs showed that vitamin D supplementation (OR: 0.72; 95% CI: 0.56-0.92) is associated with a reduced risk of wheeze/asthma. A positive trend for omega-3 fatty acids was observed for asthma/wheeze, but this did not reach statistical significance (OR: 0.70; 95% CI: 0.45-1.08). Omega-3 supplementation was also associated with a non-significant decreased risk of allergic rhinitis (OR: 0.76; 95% CI: 0.56-1.04). Neither vitamin D nor omega-3 fatty acids were associated with an altered risk of AD or food allergy.

Conclusions: Prenatal supplementation with vitamin D may have beneficial effects for prevention of asthma. Additional nutritional factors seem to be required for modulating the risk of skin and gastrointestinal outcomes. We found no consistent evidence regarding other dietary factors, perhaps due to differences in study design and host features that were not considered. While confirmatory studies are required, there is also a need for performing RCTs beyond single nutrients/foods.

KEYWORDS

AD, allergic diseases, allergic rhinitis, allergic rhinoconjunctivitis, asthma, children, eczema, food allergy, hay fever, infants, maternal diet, pregnancy, prevention, seasonal allergies, wheeze

1 | INTRODUCTION

Allergic diseases are an increasing public health concern.¹ The most common allergic conditions are asthma, allergic rhinitis/hay fever/

seasonal allergies, eczema/atopic dermatitis (AD) (from now on referred to as eczema), and food allergies. Worldwide, an estimated 300 million people suffer from asthma, and an additional 400 million have allergic rhinitis.² In addition, 570 million children and 116

million adults suffer from eczema, and 200-250 million people suffer from food allergies.³ These allergic diseases may often co-exist in one individual.⁴ Allergies can affect quality of life and reduce academic performance and income,⁵ and in rare cases can be fatal.^{6,7}

Maternal diet in pregnancy is an important, potentially modifiable factor in allergy prevention. Several studies have investigated the role of prenatal diet on offspring allergy, but the findings are conflicting, and there is a paucity of randomized controlled trials (RCTs). An evidence synthesis is paramount in order to adequately appreciate the underlying evidence.

A number of review papers have been published focusing on maternal diet during pregnancy, lactation, or early life⁸⁻¹¹ and allergy prevention.⁸⁻¹² Our review follows and extends a recent systematic review and meta-analysis by Garcia-Larsen et al,¹³ who summarized the evidence up to December 2017. Garcia-Larsen et al¹³ considered dietary intake not only during pregnancy, but also in lactation and infancy; their main conclusions¹³ were that maternal omega-3 fatty acid supplementation during pregnancy and lactation may reduce the risk of allergic sensitization to egg and peanut. However, only three of the interventional studies regarding omega-3 fatty acids included in that review were conducted in pregnancy only. In addition, the review¹³ showed that maternal probiotic supplementation in the postnatal period reduced the risk of offspring eczema and allergic sensitization to cow's milk, and none of the probiotic studies were limited to pregnancy. Since the publication of that systematic review, new studies¹⁴⁻⁴⁰ evaluating maternal diet during pregnancy and childhood allergic outcomes have been published. We reviewed diet during pregnancy only as not all mothers breastfeed or introduce solids to their infants in a similar fashion. We therefore reviewed literature regarding the pregnancy period only, to investigate specific pregnancy-related dietary associations.

Globally, recommendations on dietary intake during pregnancy for prevention of allergic disease are sparse. The Australasian guideline recommends eating fatty fish or taking omega-3 fatty acid supplements during pregnancy to reduce eczema in the offspring.⁴¹ Other guidelines from the UK,⁴² Europe,⁴³ and the United States⁴⁴ only give recommendations pertaining to food allergen avoidance in pregnancy. They state that food allergens should not be avoided during pregnancy, but give no further dietary recommendations for allergy prevention. The German allergy prevention guidelines⁴⁵ recommend that pregnant women should follow a healthy diet, need not avoid food allergens, and should eat fish regularly. In support of this, the UK prevention guidelines state that omega-3 fatty acids (found in oily fish such as salmon, trout, mackerel, and fresh [not canned] tuna) during pregnancy may help reduce the risk of eczema and allergic sensitization (development of antibodies to allergens) in early life.

The goal of this review was to summarize studies which examined the impact of maternal diet during pregnancy on offspring's allergic outcomes. We also compared the amount of nutrients or foods consumed and food patterns identified, which have been shown to have an allergy preventative effect to current US dietary guidelines.

Key Message

The systematic review focused on diet during pregnancy only, as not all mothers breastfeed or introduce solids to their infants in a similar fashion. Thus, we hence provide here pregnancy-specific dietary recommendations. Based on RCTs, we found that prenatal supplementation with vitamin D in doses higher than recommended by most countries may have beneficial effects for prevention of asthma in the offspring. Doses used in RCTs were 800 IU, 2400 IU, and 4000 IU. The European Food Safety Authority (EFSA) (<https://www.efsa.europa.eu/en/press/news/161028>) and the Institute of Medicine (<https://www.aafp.org/news/health-of-the-public/20101201omrpt-vitdcal.html>) recommend 600 IU per day for pregnant women. Summarizing the data from observational studies, we found that the following nutrients, foods, and diet patterns appear to be associated with reduced allergic outcomes: vitamin A (allergic rhinitis), vitamin D (asthma, allergic rhinitis, and sensitization), copper (food allergy, eczema, and allergic sensitization), vitamin C (food allergy and allergic sensitization), vitamin E (eczema and asthma), calcium (eczema), zinc (eczema and asthma), beta-carotene (eczema), magnesium (eczema), dairy/probiotic-containing foods (eczema and allergic rhinitis), Mediterranean and Western diet (asthma/wheeze), Mediterranean and soy/fish/nuts diet (allergic sensitization), fruit and vegetables (asthma and eczema), fish (sensitization), nuts (asthma and allergic rhinitis), fish/fatty fish (asthma), and meat (asthma). However, the doses of foods or nutrients studied were heterogeneous and we were unable to relate these to current nutrition guidelines. We found no consistent evidence regarding other dietary factors, perhaps due to differences in study design and host features such as nutrient intake/levels at the start of the study that were not considered. There is a need for confirmatory studies and RCTs beyond single nutrients.

2 | METHODS

2.1 | Studies included in this review

2.1.1 | Types of studies

We included RCTs, observational (cross-sectional and cohort), and case-control studies, which (a) considered the time period of pregnancy alone, (b) reported allergic outcomes or allergic sensitization status in the offspring, and (c) were written in English. Studies examining food allergen intake were excluded. Studies that included intake of major allergens such as milk or nuts were included, as long as the outcome did not relate to the particular food allergen. As an example, studies on tree nut intake were included only when an association

with an allergy outcome other than a tree nut was studied, such as asthma.⁴⁶ The latest data indicate that early introduction of a food allergen specifically prevents food allergy to that allergen only.⁴⁷ We were therefore confident that any outcome noted was less likely to be related to the nutritional composition, rather than the allergen content of the food. We also excluded case studies, case series, systematic reviews and meta-analyses, expert reviews and opinions, and editorials.

2.1.2 | Types of publications

A consensus was reached on including full-text publications only, as there were discrepancies in some of the data published in abstract form (as presented at conferences) when compared to the full paper publications.

2.1.3 | Types of participants

Pregnant women were the target group for this systematic review, provided there was allergic outcome data published for their offspring. Both general population and populations at high risk of atopy were included.

2.1.4 | Types of interventions in RCT

Studies that used supplementation or specific foods (eg, fatty fish) during pregnancy alone, irrespective of dose, formulation, or mode of delivery and composition (eg, oil and tablet), were included. Trials were not included if the intervention(s) had been continued beyond pregnancy through breastfeeding or given only to the infant.

2.1.5 | Types of outcome measures

Studies were included if they reported on allergic outcomes or allergic sensitization status in the offspring, either as a primary end-point or as a secondary end-point. Allergic outcomes were defined as wheeze/asthma, allergic rhinitis/rhinoconjunctivitis/hay fever, food allergy, eczema, allergic sensitization measured by skin prick tests/specific IgE tests to food/aero/any allergen, or total IgE levels. Please see Supplementary File S1 for a list of the search terms used in respective databases. An expert panel from the United States, Europe, and Australia commented on the search strategy and the list of identified and included studies. Additional studies were located through searching the references cited in identified studies and systematic reviews, and through discussion with the expert panel.

2.2 | Dietary guidelines for food/nutrient intake in pregnancy

To compare our study findings with current dietary guidelines, we used the following approach to calculate food or nutrient intake: For

micronutrient comparisons, the US dietary reference intakes (DRI) and the United States Department of Agriculture (USDA) www.choosemyplate.gov guidelines⁴⁸ for pregnant women were used. Macronutrient requirements and number of portions for the second trimester of pregnancy were calculated, assuming a daily caloric intake of 2400 kcal, comprising 71 g of protein, 53.33-93.33g of total fat, <27 g of saturated fat, 13 g of linoleic acid (LA), 1.4 g of alpha-linolenic acid (AI), 175 g of carbohydrates, 28 g of fiber, <12 g of added sugar, 2 cups (2× approx. 125 g) of fruit, 3 cups (3 × 125 g) of vegetables, 8 oz of grains (8 × 30 g), 6.5 oz of high-protein foods (6.5 × 30 g), and 3 cups (3 × 200-250 mL) of dairy. Recommended fish intake was assumed to be 2-3 servings (in total approx. 8-12 oz/240-360 g) per week as per the United States Food and Drug Administration (FDA) (Best Choices: <https://www.fda.gov/food/consumers/eating-fish-what-pregnant-women-and-parents-should-know>).

2.3 | Data collection

2.3.1 | Ethics application/Institutional Review Board

This study is considered exempt from Institutional Review Board review by the University of Colorado School of Medicine Combined Institutional Review Board.

2.4 | Study selection

The titles and abstracts of articles considered for inclusion were independently assessed by MBA, MP, and AM and categorized as included, not included, or potentially relevant. CV then checked each assessment. For studies considered potentially relevant, full-text copies were obtained, and inclusion of the studies was discussed by CV, MBA, MP, and AM. Any discrepancies were resolved either by consensus among the authors, or, if there was lack of consensus, by discussion with the expert panel. All the studies in this publication were reviewed and approved by both the authors and the expert panel.

2.5 | Risk of bias assessment

Risk of bias assessment was undertaken independently by MBA and CV using a similar approach as in Garcia-Larsen et al¹³ As in the paper by Garcia-Larsen-et al,¹³ we used modified versions of the Cochrane Collaboration Risk of Bias tool for intervention trials and the National Institute for Clinical Excellence methodological checklist for cohort and case-control studies.⁴⁹ For intervention trials, risk of bias assessment included assessment of: (a) selection bias, including sequence generation and allocation concealment; (b) assessment bias, including blinding of outcome assessors and ensuring outcomes were determined by validated assessment tools; and (c) attrition bias, which was considered high when outcome data were reported in <70% of randomized participants. Risk of bias for cohort

and case-control studies included assessment of: (a) selection bias, which was considered low if cases and controls were recruited from similar populations and had a similar attrition rate <20%; (b) assessment bias, which included blinding of outcome assessors and use of validated assessment tools; and (c) confounding bias (did study design and analysis account for relevant confounders?). Conflicts of interest were noted if industry was involved in any aspect of the study or if authors received funding for other activities from relevant industry partners. Using these components, each study was graded as high, medium, or low. Supplementary File S2 includes the grading for risk of bias assessment. In order to minimize publication bias, we performed a comprehensive search of the literature and included experts in the field to ensure that no relevant study was missed. We did not include gray literature (unpublished information or information that is not produced by commercial publishers) in order to avoid discrepancies with published data. None of the meta-analyses included more than 10 papers, and we therefore did not use funnel plot and Egger tests to formally test for publication bias.

2.6 | Data extraction and reporting

Data were extracted by three authors (AM, MBA, and MP). New articles were added into the excel sheets downloaded from Garcia-Larsen et al¹³. Inclusion and exclusion criteria differed between the current review and that of Garcia-Larsen et al¹³; some entries were edited, modified, or deleted. The edited files were reviewed for quality and completeness by CV. Detailed information on study characteristics was recorded by CV, MP, and MBA and appears in Supplementary File S3. Any discrepancies regarding the classification of dietary measures and offspring allergic outcomes were discussed and resolved within the group of authors. Classification of maternal dietary measures and offspring allergic outcomes were double-checked by the expert panel. The authors' names, institutions, journals, or data were not blinded during data extraction.

2.7 | Data analysis

2.7.1 | Observational studies

For wheeze, studies were split into two categories: infants and young children (0- <3 years) and older children (3 years and older). All studies reporting on allergic rhinitis included offspring with ages over 5 years. All ages were included for food allergy, eczema, and allergic sensitization.

The reported data were plotted. If multiple odds ratios (ORs) were reported for a single study, only the highest exposure compared to the lowest exposure OR was used. Meta-analysis of observational studies was not appropriate because the studies were heterogeneous in focus, design, and target populations. We used forest plots to visually summarize all significant outcomes found by the studies, that is, the adjusted OR, adjusted hazard ratio (HR), or adjusted risk ratio

(RR), and their corresponding 95% confidence intervals. If more than one significant outcome was found in a manuscript, then only the largest observed difference was included in this review. For example, if authors compared quartiles, by comparing Q1 vs Q2 and Q1 vs Q4 for one nutrient in the same age-group, only the largest difference would have been included. We understand that it is better to report the estimates regardless of their statistical significance in order to give readers an appreciation of the magnitude of the effect sizes and that statistical significance is purely a function of the sample size of a study and hence provides information only of the random error. It does not give any indication of the magnitude of the estimates or potential systematic errors. However, our goal here was to provide a visual presentation of the data, and the Supplementary Files S1-S9 contain the full set of information. Therefore, findings from observational studies were synthesized narratively by grouping studies according to allergy outcome and nutrient, food, or diet pattern studied: nutrients, including vitamins and minerals, omega-3 fatty acids and fatty fish, fruit and vegetables, other foods, and food patterns.

2.7.2 | Randomized clinical trial meta-analyses

For each exposure-disease outcome pair included in the RCTs, separate meta-analyses were performed. Because the number of studies was limited, several meta-analyses included fewer than four data points. All meta-analyses included at least two studies.⁵⁰ DerSimonian-Laird random-effects models were used to perform the meta-analyses and estimate the pooled ORs using the *rmeta* (v 3.0)⁵¹ package in R. The heterogeneity of the studies was calculated using the Cochran χ^2 and I^2 statistics. Forest plots were created for each of the diseases using the *forestplot* (v1.9)⁵² package in R (v3.5.1). Where studies reported outcomes in the same children at more than one time point, we included each outcome in the meta-analysis as well as the combined outcomes. As the different analyses showed no changes in the outcomes, the overall decision was to use food allergy outcomes at 3 years; eczema at 3 years; allergic sensitization at 3 years; and asthma, rhinitis, and allergic sensitization at 6 years.

3 | RESULTS

3.1 | Results of literature search and characteristics of included studies

We included all the relevant published studies on the topic of maternal dietary intake during pregnancy alone identified by Garcia-Larsen et al¹³ (n = 68 papers).^{14-40,46,53-119} An additional search was performed using the same search terms to include studies published after, or potentially missed by their search, including observational studies from August 2013 to February 2019 and intervention studies from January 2017 to February 2019 (Figure 1). Based on this additional search, we identified an additional 27 publications. Of them, 25 were cohort or case-control studies¹⁴⁻³⁸ and 2 were RCTs.^{39,40} In




FIGURE 1 Flow chart for study inclusion

total, our systematic review is based on 95 papers (17 RCTs and 78 cohort and case-control studies). The numbers of pregnant women and their offspring studied at different time points and presented in different publications are shown in the supplementary Files S1-S9. Supplementary File S4 has information on nutrients, including vitamins and minerals; Supplementary File S5 contains information on fats and fatty fish; Supplementary File S6 is on fruit and vegetables; Supplementary File S7 contains information on other foods; and Supplementary File S8 is on food patterns.

Sufficient numbers (at least two) of RCTs were found for vitamin D, vitamins C and/or E, and omega-3 fatty acids. We performed a total of 9 meta-analyses.

3.2 | Quality and risk of bias assessment


The 95 papers were generated from 50 studies. The 50 studies included 11 RCTs^{39,40,53-67} (7/11 normal risk), 2 cross-sectional^{21,74} (both normal) and 30 cohort studies^{15,18-20,22-30,32-37,46,68-70,72,73,75-78,80-108,110,112,115-119} (all normal risk), 3 case-control designs nested within cohorts^{16,17,31,38,79,113,114} (all normal risk), and 4 case-control studies^{14,71,109,111} (3 normal and 1

high risk). Due to low numbers of RCTs that could be included in the meta-analysis and low numbers of studies including high-risk populations in the observational studies, these were not separated. Of the studies, 8 (7 observational and 1 RCT) (16%) were considered to be of high quality, 10 (8 observational and 2 RCTs) (20%) were of uncertain quality, and 32 (24 observational and 8 RCTs) (64%) were of low quality. The results for the quality assessment of the studies are shown in Supplementary File S2.


3.3 | Prevention strategies in pregnant women

Supplementary excel Files S4-S8 summarize the results of all the 95 publications. The excel files contain, if available for each paper, the last name of the first author, publication year, study design, dietary exposure, dietary intake time, outcome type, offspring's age at assessment, and every relevant point estimate reported with its associated 95% confidence intervals and p-values, as well as the relevant comparison of interest that each point estimate describes. Due to the heterogeneity between papers on the measurement of dietary intake, the analysis of the data, and the reporting of the results, data from observational studies were not pooled in meta-analyses.

(A) Anti-oxidants: Vitamin C and Vitamin E



(B) Omega-3 fatty acids *



(C) Vitamin D




FIGURE 2 A-C, Randomized controlled trials with asthma/wheeze as an outcome

3.4 | Dietary intake

Dietary associations measured in observational studies depend on the validity of the measures used to study dietary intake.¹²⁰ This important methodological factor has so far been omitted from past systematic reviews. Methods for measuring dietary intake differ greatly between studies. We have summarized the approaches for

assessing dietary intake and adherence to study interventions used in the RCTs in Supplementary File S3. Of the 17 publications reporting on RCT outcomes, 8 papers^{39,40,59-62,65,66} mentioned that they monitored adherence and 6 papers^{54,55,57,58,63,64} did not monitor intake. For the observational studies, instruments used to measure food or nutrient intake were validated food frequency questionnaires (FFQs),^{15,18,21,23,25-27,30,33,34,36,46,70,75-78,80-84,87-107,110,115,118}

TABLE 1 Correct ref numbers

Food and dietary patterns	Asthma/wheeze	Hayfever/rhinitis	Eczema	Food allergy	Sensitization
Total	62 papers ^{14-16,19-24,28,30-33,35-38,46,68-70,74-80,82-93,95,96,98-101,103-111,113,117-119}	19 papers ^{15,16,18,21,24,25,31,32,46,70,90-93,103,104,106,107,113}	38 papers ^{15,16,19,20,24,26,29,31,37,68,70,72,76,77,81,82,84-88,90,94-101,103,104,107,108,112,113,116,118}	5 papers ^{34,71,77,115,118}	19 papers ^{16-18,24,27,31,68,73,75-77,82,86,102,110,112-114,118}
Fruit and/or vegetables (Supplementary excel file 6 Fruits and Vegetables)	11 papers ^{20,28,36,69,75,78,90,99,106,108} Fruit, vegetables, leafy vegetables, roots and potatoes, citrus fruits, Malaeous fruits, berries, fruit and berry juices, Green and yellow vegetables, Folate vegetables, apples	1 paper ¹⁰⁶ classified as fruits, vegetables, leafy and root vegetables, berries, malicious, citrus fruit and juices	4 papers ^{20,99,108,112} fruits, vegetables, citrus fruits, apples, green leafy vegetables, green and yellow vegetables, carrots, spinach, cabbage, celery, tomato, sweet pepper, salad, and juice	1 papers ¹¹⁵ fruit and berries, vegetables and roots	3 papers ^{77,102,112} fruits, vegetables, malicious fruits, citrus fruits, berries, apples, bananas, strawberry, exotic fruit, root vegetables, carrots, spinach, cabbage, celery, tomatoes, sweet pepper, salad, and fruit and vegetable juices
Fats and fatty fish (Supplementary excel file 5 Fish and Fats)	12 papers ^{20,32,36,74,78,85,95,96,104-106,111}	432,90,104,106	9 ^{20,29,85,95-97,104,112,116}	1	4 papers ^{27,73,102,112}
Fat and fatty acid intake (4 papers), classified as total fat intake, omega 3 or omega 6 poly-unsaturated fatty acids or other types of fatty acids, and specific fatty acids (we also included fish intake and fish oil supplementation in this group);	33 papers ^{14,15,22,23,25,26,28-30,33,35,36,38,68,77,79,83,84,86-89,91,98-100,107,110,117-119} multivitamin, pre-natal vitamins, folic acid, vitamin A, vitamin C, vitamin D, vitamin E, vitamin K, B-vitamins and their precursors, zinc, calcium, selenium, copper, magnesium, and manganese	6 papers ^{15,18,25,103,107} folate, riboflavin, retinol, vitamin A, vitamin D, vitamin E, vitamin K, and their precursors, zinc, copper, selenium, magnesium, calcium, manganese	14 papers ^{15,26,68,77,84,87,88,98-100,103,107,116,118} vitamins A, C, E, and their precursors, vitamin D, vitamin B and folic acid, mineral intake zinc, calcium, copper, magnesium, manganese and selenium	4 papers ^{7,71,115,118} antioxidant vitamins C, E, and B-carotene, vitamin D and folic acid, mineral intake, copper and zinc	9 papers ^{7,18,68,77,86,102,110,114,118} Vitamin A, Vitamin C, Vitamin D, Vitamin E, Folic acids, Prenatal vitamins, Multivitamins, Anti-oxidants and their precursors, Zinc, Selenium
Other dietary exposures (Supplementary excel file 7 other foods)	25 papers ^{16,19,20,24,31,32,37,46,69,70,75,76,80,85,92,93,95,96,98,105,106,108,109,111,117} sugar, milk/dairy products, grains (pasta), nuts, meat, fish, chocolate candy, soft drinks/artificial sweeteners, egg, alcohol, glucose and dietary water	8 papers ^{16,31,32,46,70,92,93,106} sugar, milk/dairy products, cereals/grains, fish, seafood, meat, chocolate candy, nuts, soft drinks/artificial sweetener, alcohol.	22 papers ^{16,19,20,26,29,31,37,70,72,76,81,85,90,94-98,108,112,116} alcohol, probiotic drinks, fast foods, meat, dairy, cereals, fish and shellfish, chocolate, cheese, natto, eggs, cream, seeds and nuts.	1 papers ¹¹⁵ Cereal and dairy	7 papers ^{16,27,31,73,76,102,112} sugar, fish, nuts, legumes, seeds, mild/dairy products, cereals/grains, egg, chocolate candy, juices and alcohol
Food patterns (Supplementary excel file 8 dietary patterns)	8 papers ^{20,21,24,74,75,82,101,113} Mediterranean, Health conscious, Vegetarian, Prudent, Western, Confectionery, Traditional, Japanese, Vegetable, Fruit and white Rice, Seafood and Noodles Pasta, Cheese and Processed meat, Alternative Healthy Eating index	3 papers ^{21,24,113} Mediterranean, Health conscious, Traditional, Processed, Vegetarian, Confectionery, Japanese, Vegetable, Fruit and white Rice, Noodles Pasta, Cheese and Processed meat, Alternative Healthy Eating index	5 ^{20,24,82,101,113} Mediterranean, Health conscious, Vegetarian, Prudent, Western, Confectionery, Traditional, Japanese, Vegetable, Fruit and white Rice, Seafood and Noodles Pasta, Cheese and Processed meat, Alternative Healthy Eating index	0	5 papers ^{17,24,27,75,82,102,112,113} Mediterranean diet, Alternate Healthy Eating Index modified for Pregnancy score, Prudent diet, Western diet, Vegetable, Fruit and white Rice, Seafood and Noodles, Pasta, Cheese and Processed meat, Vegetable, Fruit and white Rice, Seafood and Noodles, Pasta, Cheese and Processed meat, and white Rice, Seafood and Noodles, Pasta, Cheese and Processed meat, Healthy Conscious, Traditional, Confectionery, Traditional and Vegetarian

unvalidated FFQs,^{16,17,19,28,29,31,32,73,111-114,119} questionnaires,^{14,20,22,35,38,68,69,71,72,74,79,85,86,108,116} interviews,²⁴ information obtained from medical records,¹⁰⁹ 3/5-day food records and/or 24-hour recalls and questionnaires^{37,117} These points are important to take into account when reading the data.

3.5 | Asthma/wheeze outcomes

Early childhood wheeze may not lead to allergic asthma outcomes in later childhood.^{121,122} To distinguish between transient wheeze/asthma in infancy and early childhood, and later wheeze/asthma, which may be a stronger predictor of subsequent allergic asthma outcomes, we have separately reported observational study manuscripts on maternal diet and wheeze/asthma into those which considered offspring wheeze/asthma between 0 and <3 years and for offspring 3 years and older. For the meta-analysis, we have included all ages. Allergic outcomes and how these were measured are shown in supplementary File S3.

3.5.1 | Randomized controlled trials

We had a limited number of studies to perform meta-analysis of anti-oxidant (vitamins C and E; 2 studies—children of 1-2 years),^{58,61} vitamin D (3 studies—children of 3 years old),^{55,57,60} and omega-3 fatty acid (4 studies; 5 papers; children of age 6m, 5 - 16 years)^{62,63,66,67,72} intake or supplementation against wheeze/asthma outcomes in the offspring (Figure 2A-C). Our meta-analyses indicate that vitamin D during pregnancy had a statistically significant protective effect on wheeze/asthma in the offspring (OR: 0.72; 95% CI: 0.56 - 0.92). Although there was a trend toward a protective effect, maternal supplementation/intake during pregnancy with omega-3 fatty acid (OR: 0.70; 95% CI: 0.45-1.08) did not reach statistical significance on wheeze/asthma outcomes in the offspring. If we remove the study by Noakes et al who reported on wheeze in the first six months of life, significance is still

not reached (OR: 0.63; 95% CI: 0.4-1.00—meta-analysis not shown). Using data from children at 1-2 years, anti-oxidant did not show a protective effect against asthma/wheeze (OR: 0.67; 95% CI: 0.29-1.56).

3.5.2 | Observational studies

Many different outcomes have been reported as summarized in supplementary File S9. Sixty-two papers from observational studies^{14-16,19-26,28,30-33,35-38,46,68-70,74-80,82-93,95,96,98-101,103-111,113,117-119} reported on the association between maternal diet during pregnancy and asthma/wheeze as an outcome (Table 1; Figures 3A and B and 4A and B). We summarized all the significant outcomes reported in Figure 3 to give a visual presentation of the outcomes rather than making a statistical conclusion. Figure 4 indicates that a large number of foods/nutrients/food patterns were associated with an increased or reduced risk of asthma/wheeze in children 3 years and over. The most significant protective factors were foods associated with a Mediterranean diet, meat, vitamin D, and fatty fish. The most significant factors associated with an increased risk were pasta, fish sticks, and arachidonic acid. From the summarized data, no firm conclusion can be drawn in regard to the protective effect of nutrients/foods/food patterns studied on allergic disease. However, when considering asthma, reduced offspring allergic outcomes were associated with intake of copper, vitamin D, Mediterranean diet and Western diet, fish/fatty fish, tree nuts, fruit, and meat (type of meat not specified) in the observational studies included.

3.6 | Allergic rhinitis outcomes

Figures 5 and 6 summarize allergic rhinitis outcomes in the offspring for both observational and randomized controlled studies, respectively. All observational studies with significant outcomes are shown in Figure 6A and B. The studies shown in both these figures included results for offspring aged 3 years and above.




FIGURE 3 A and B, Observational studies with asthma/wheeze (age 0-3 y) as an outcome





FIGURE 4 A and B, Observational studies with asthma/wheeze (age 3 y and above) as an outcome



3.6.1 | Randomized controlled trials

We were able to perform meta-analysis of omega-3 fatty acid supplementation (2 studies, 3 papers—all children over 5 years) ^{39,54,66,67} against allergic rhinitis outcomes in the offspring

(Figure 5). Although we found a potential protective effect based on the meta-analysis combining two studies of omega-3 supplementation during pregnancy on allergic rhinitis in the offspring, the effect showed a trend toward statistical significance (OR: 0.76; 95% CI: 0.56-1.04).

3.6.2 | Observational studies

Nineteen papers from observational studies^{15,16,18,21,24,25,31,32,46,70,90-93,103,104,106,107,113} reported an association between maternal diet during pregnancy and offspring allergic rhinitis/rhinoconjunctivitis as an outcome. Different studies reported different terms to describe outcomes and definitions (ie, some studies referred to ever suffering from hay fever and others reported on current allergic rhinitis) were not always similar, though the same disease group was being studied. The results are described and summarized in both Table 1 and Figure 6.

Figure 6A and B indicates that vitamin D, vitamin A, fatty fish, polyunsaturated fatty acids (PUFA), alpha-linolenic acid, peanut, pistachio, and probiotic milk products were associated with reduced risk of allergic rhinitis. Fish (unspecified type), the ratio of omega-6 to omega-3 fatty acids, butter, and fruit were all associated with increased risk of allergic rhinitis. As above, due to heterogeneity between studies, these data were summarized narratively and not pooled for meta-analysis. Vitamin A, vitamin D, dairy, nuts, and probiotic-containing foods showed associations with reduced allergic rhinitis outcomes.

3.7 | Eczema outcomes

Eczema usually develops in infancy¹²³ prior to food allergy and other allergic manifestations, making it a target for allergy prevention studies. Data presented in this systematic review include offspring data from 6 months onwards.

3.7.1 | Randomized controlled trials

We were able to perform meta-analysis on the effects of vitamin D^{55,57,60} (3 studies) and omega-3 fatty acid intake⁶² or supplementation^{39,53,54,56,65-67} (4 studies, 8 papers; 4 used in meta-analysis) on offspring eczema outcomes (Figure 7A,B). None of the dietary exposures during pregnancy had a significant effect on offspring eczema (vitamin D [OR: 0.90; 95% CI: 0.71 - 1.15] and omega-3 [OR: 1.07; 95% CI: 0.82-1.40]).

3.7.2 | Observational studies

Thirty-eight papers from observational studies^{15,16,19,20,24,26,29,31,37,68,70,72,76,77,81,82,84-88,90,94-101,103,104,107,108,112,113,116,118} reported the association between maternal diet during pregnancy and eczema in their offspring (Table 1; Figure 8). Due to heterogeneity between studies, these data were not pooled for meta-analysis.

Figure 8A and B summarize the significant associations published between various components of maternal diet during pregnancy and offspring eczema that were reported from observational studies. In

terms of vitamins and minerals, studies presented significant associations for a decreased risk of offspring eczema with maternal intake of beta-carotene, vitamin E, zinc, calcium, magnesium, and copper during pregnancy. Paradoxically, studies showed both positive and protective associations between maternal intake of vitamins C and D and offspring eczema. Intake of fatty fish, margarine, vegetable oil, total fat, omega-6 fatty acid, omega-3/omega-6 ratio, linoleic acid, alpha-linolenic acid, natto, and MUFAs all was positively associated with offspring risk of eczema. Cholesterol and arachidonic acid showed a decreased association with eczema. Looking at other foods, fast foods, shellfish, alcohol, and meat (not defined) were associated with an increased risk of offspring dermatitis. Maternal intake of milk, fish, vegetables, and apples was associated with a reduced risk of developing AD. Intake of copper, vitamin E, calcium, zinc, beta-carotene, magnesium, dairy/probiotic foods, vegetables, and fruit was associated with reduced eczema outcomes.

3.8 | Food allergy outcomes

3.8.1 | Randomized controlled trials

There were two (three papers)^{56,65,66} RCTs that studied the effect of maternal intake of omega-3 fatty acid during pregnancy and food allergy outcomes in the offspring. The meta-analysis is shown in Figure 9, revealing that maternal intake/supplementation of omega-3 fatty acids during pregnancy had no significant effect on food allergy in offspring (OR: 1.18; 95% CI: 0.56 - 2.46).

3.8.2 | Observational studies

Five publications from observational studies^{34,71,77,115,118} discussed associations between maternal diet during pregnancy and food allergy in their offspring (Table 1). Figure 10 summarizes significant results from observational studies regarding the association between various components of maternal diet during pregnancy and food allergy in offspring. The studies reported protective associations between maternal intake of copper and vitamin C and risk of food allergy outcomes in the offspring. Maternal intake of vegetables and vitamin D was associated with increased risk of food allergy in the offspring.

3.9 | Allergic sensitization outcomes

Allergic sensitization in the included studies measured both allergic sensitization to food and/or aero-allergens, using skin prick test (SPT) and/or specific IgE levels. Different outcome measures were used, including increased total IgE, a range of specific IgE or SPT to foods and aero-allergens, and amalgamation using the term atopy to indicate any level of allergic sensitization.

Omega-3 fatty acids





FIGURE 5 Randomized controlled trials with allergic rhinitis/hay fever as an outcome

(A)



(B)





FIGURE 6 A and B, Observational studies with allergic rhinitis/hay fever as an outcome

3.9.1 | Randomized controlled studies

Three^{55,57,60} RCTs reported results on the effects of vitamin D on allergic sensitization in the offspring, and five reported on the effects of omega-3 fatty acid intake⁶² or supplementation^{39,54,56,65-67}

on the same outcome. The meta-analysis is shown in Figure 11A, B. The meta-analyses showed no effects for either maternal intake of vitamin D (OR: 0.91; 95% CI: 0.68 - 1.21) or intake/supplementation of omega-3 fatty acid (OR: 0.102; 95% CI: 0.77 - 1.36) during pregnancy with offspring allergic sensitization outcomes.

(A) Vitamin D



(B) Omega-3 fatty acids




FIGURE 7 A and B, Randomized controlled trials with atopic dermatitis as an outcome

3.9.2 | Observational studies

Nineteen papers from observational studies ^{16-18,24,27,31,68,73,75-77,82,86,102,110,112-114,118} studied the association between maternal diet during pregnancy and allergic sensitization as an outcome.

Data shown in Figure 12 and B and summarized in Table 1 indicate that vitamin C, vitamin D, copper, omega-6 PUFAs, PUFA, butter, fish, the Mediterranean diet, and the seafood and noodle diet pattern were associated with reduced allergic sensitization. Fruit, celery, sweet pepper, free sugar, fatty fish, vegetable fat, and vegetarian and health-conscious diet were associated with increased prevalence of allergic sensitization. Intake of vitamin D, copper, vitamin C, and diet containing soy, fish, and nuts was associated with reduced prevalence of sensitization.

3.10 | Dietary intake and its association with allergy outcomes

Where data allowed, we summarized the dose of nutrients/food intake associated with allergy outcomes in observational studies

(Tables 2 and 3). We find no clear association between dose of nutrient/food required to reduce or increase any specific allergy outcomes. Of note is that the dose/amount showing an association frequently does not comply with national dietary recommendations from the United States.¹²⁴ The following associations between diet pattern, food and/or nutrient intake and allergic outcomes was reported in the observational studies: Vitamin A (allergic rhinitis), vitamin D (asthma, allergic rhinitis, and sensitization), copper (food allergy, eczema, and allergic sensitization), vitamin C (food allergy and allergic sensitization), vitamin E (eczema and asthma), calcium (eczema), zinc (eczema and asthma), beta-carotene (eczema), magnesium (eczema), dairy/probiotic-containing foods (eczema and allergic rhinitis), Mediterranean diet and Western diet (asthma/ wheeze), Mediterranean and soy/fish/nuts diet (allergic sensitization), fruit and vegetables (asthma and eczema), fish (sensitization), nuts (asthma and allergic rhinitis), and meat (asthma). We suggest that researchers use nutrition guidelines to set predetermined cutoffs for food intake to (a) compare outcomes in observational studies and (b) determine dose given during RCTs. This will help ensure safe dietary interventions and will also allow comparison of future studies.

(A)





FIGURE 8 A and B, Observational studies with atopic dermatitis as an outcome

(B)



4 | DISCUSSION

4.1 | Summary of findings

We find insufficient evidence to provide guidance on diet diversity (no published reports), diet patterns, diet indices or specific foods, food groups, and macro- or micronutrients that should be consumed

or avoided during pregnancy for the prevention of allergic diseases in the offspring. However, based on 3 RCTs, we did find that intake of vitamin D, higher than EFSA and IOM recommendations, was significantly associated with a reduced risk of wheeze/asthma. This needs to be further studied before guidelines are changed, particularly as intake of vitamin D in both lactation¹²⁵ and early life is associated with increased food allergy in the offspring.

Omega-3 fatty acids




FIGURE 9 Randomized controlled trials with food allergy as an outcome




FIGURE 10 Observational studies with food allergy as an outcome

4.2 | Strengths and limitations

We included the most up-to-date studies on dietary intake during pregnancy and prevention of allergic disease in their infants. Studies from Europe, North America, Asia, and Australia were included. We used the stringent standard employed by Garcia-Larsen et al¹³ to conduct the search and perform the risk of bias assessment and data extraction. We brought together a panel of international experts to review the data and conclusions of this review.

The review was limited by the quality of studies included. Some 64% of the studies included were deemed to be of low quality. The pool of data is also relatively small with 50 studies contributing to 95 papers. The observational studies were highly heterogeneous and were therefore not considered to be suitable for meta-analysis. There is a paucity of RCTs in this field; only four meta-analyses were performed using data from RCTs. Finally, none of the studies looked at diet diversity during pregnancy, this may be partly due to the uncertainties in defining diet diversity, which has now been addressed in an EAACI position paper.⁹

4.3 | Comparison of findings to previous studies


The systematic review by Vadhaninia et al¹²⁶ based on RCTs assessed the impact of omega-3 fatty acid supplementation solely in pregnancy and found that based on pooled estimates, omega-3 fatty acid intake significantly reduces allergic sensitization to egg and

peanut in the offspring. We did not find a significant effect on any allergic sensitization outcome. The discrepancy may be attributed to the fact that one of the studies included by Vadhaninia et al¹²⁶ was excluded from our review, as maternal supplementation continued up to 2.5 months post-delivery.¹²⁷ Our findings are also different from the conclusions drawn by Garcia-Larsen et al.¹³ The difference probably lies in the choice of studies included in each systematic review. Garcia-Larsen et al¹³ included studies in pregnancy, lactation, and early life. We considered studies in pregnancy alone. It is therefore possible that supplementation with omega-3 fatty acid, which continues during breastfeeding and early life, may be more effective than just during pregnancy alone to prevent allergic outcomes in the offspring.

In terms of probiotic supplementation, published literature suggests that probiotic supplementation during late pregnancy, breastfeeding, and early life^{13,43,128-135} may be associated with reduced risk of the development of eczema. Substantial uncertainties remain regarding the specific strain required and the optimal timing of supplementation. We were unable to find any RCTs using probiotic supplementation in pregnancy only. We did find only one observational study that examined the association between maternal intake of probiotics and offspring allergy, but the authors found no association with allergic outcomes.¹⁰⁶

We were able to demonstrate that vitamin D supplementation in pregnancy may reduce the risk of wheeze/asthma.^{55,57,60} Vadhaninia et al¹³⁶ also concluded that prenatal supplementation of vitamin D might be associated with reduced odds of recurrent wheezing in children. We were unable to find any RCT on folic acid supplementation

(A) Vitamin D



(B) Omega-3 fatty acids




FIGURE 11 A and B, Randomized controlled trials with “any” sensitization as an outcome




FIGURE 12 A and B, Observational studies with “any” sensitization as an outcome

in pregnancy only and observational studies provided conflicting evidence.^{14,22,28,30,33,35,38,68,71,77,79,83,100,103,114,115,119} There is lack of evidence on the effect of other vitamins for the prevention of respiratory and/or allergic outcomes.

Many systematic reviews have made recommendations, which relate to maternal diet in pregnancy and allergy prevention in the offspring. The review by Lodge et al¹³⁷ did not identify any nutritional factors during pregnancy that reduced the risk of food allergy in the

TABLE 2 Comparison of food and nutrient (where available) against RDA or AI

Wheeze/asthma				
Food/nutrient intake	Reference^a	Intake	RDI or AI	% RDI or AI
Alpha-tocopherol ²⁸	Increase	6.5 mg	15 mg	43%
Calcium ⁹⁸	Reduce	571.1 mg	1000 mg	57.1%
Folate ³⁰	Increase	>578	600 µg	>96.3%
Vitamin D ¹⁵	Reduce	4.69-35 µg	15 µg	31.2%-233%
Vitamin D ¹⁵	Reduce	3.51-4.68 µg	15 µg	23.4%-31.2%
Vitamin D ⁸⁴	Reduce	724 IU (18.1 µg)	15 µg	120%
Vitamin D ²³	Reduce	4.86-17	15 µg	32.4%-113%
Vitamin D ²³	Reduce	3.44-4.85	15 µg	22.9-32.3
Vitamin D ⁹¹	Increase	8.2, 13, 16.5	15 µg	54.7%, 86.7%, 110%
Vitamin D ⁹⁸	Reduce	5.1, 6.4	15 µg	34%, 42.7%
Vitamin E ¹⁵	Reduce	6.22-7.08 mg	15 mg	41.5%, 47.2%
Vitamin E ⁸⁸	Reduce	7.07-30.30	15 mg	47.1%-202%
Vitamin E ²³	Reduce	10.3-29.4	15 mg	68.7%-98%
Vitamin E ⁹⁹	Reduce	7.3 mg	15 mg	48.6%
Zinc ⁸⁸	Reduce	14.25-30.30 mg	11 mg	126%-268%
Zinc ⁹⁹	Reduce	7, 8.5 mg	11 mg	61.9%-75.2%
Omega 6 ¹⁰⁵	Reduce	<8.03 g	13 g	61.8%
Alpha-Linolenic acid ¹⁰⁵	Increase	<7.78 g	13 g	58.9%
N-3 poly-unsaturated fatty acid ¹⁰⁵	Increase	<2.24 g	1.4 g	160%
n-3 poly-unsaturated fatty acid ⁹⁶	Reduce	2.2 g	1.4 g	157%
Alpha-Linolenic ⁹⁶	Reduce	2.3 g	1.4 g	164%
Vegetables ¹⁰⁶	Increase	<149.1	375 g	39.7%
Fruit ¹⁰⁶	Increase	326.2	250 g	130.5%
Vegetables ²⁸	Reduced	206-286 g/d	375 g	54.9%-76.3%
Free sugar ¹⁶	Increase	82.4-345.1 g/d	<12 g	687%-2875.8%
Meat ⁹⁶	Reduce	63.6 g/d	71 g	89.6%
Dairy products ⁹⁸	Reduce	280.7 g/d	250 g	115%
Allergic rhinitis				
Vitamin D - food ¹⁰⁷	Reduce	4.4-5.72 mg	15 µg	29%-38.1%
Alpha-linolenic acid ¹⁰⁴	Reduce	6.4-8.9 g	1.4 g	457%-635.7%
Fruit ¹⁰⁶	Increase	145.7-1700 g/d	250 g	58.3%-680%
Atopic dermatitis				
Beta-carotene ⁹⁹	Reduce	1923.9 and 4218.0 mg	770 µg	41.5% and 91.3%
Vitamin D ²⁶	Increase	8.6 µg/d	15 µg	57%
Vitamin E ¹⁵	Reduce	2.32-2.9 mg	15 mg	15%-0.19%
Vitamin E ⁹⁹	Reduce	86.7 mg/d	15 mg	578%
Zinc ⁸⁸	Reduce	10.81, 13.4, 22.27 mg/d	11 mg	95%, 117% 194%
Calcium ²⁶	Reduce	677.6 mg/d	1000 mg	67.8%
Omega - 6 ⁹⁶	Increase	14.1	13 g	107%
Linolenic acid ⁹⁶	Increase	10.4-11.3 g	1.4 g	743%-807%
Omega 6 ⁹⁷	Increase	10.6 g/d	13 g/d	81.5%
Total fat ⁹⁷	Increase	58.6/d	80 g/d (range 53-93 g)	3.25%

(Continues)

TABLE 2 (Continued)

Wheeze/asthma				
Food/nutrient intake	Reference ^a	Intake	RDI or AI	% RDI or AI
Alpha Linolenic acid ⁹⁷	Increase	1.9	1.4 g/d	136%
Vegetables ⁹⁹	Reduce	Total Veg 144.4 g	3 × 125 g	39%
Total dairy products ⁹⁸	Reduce	120.8 g/d	500 ml/g/d	24%
Total dairy products ²⁶	Reduce	255.3 g/d	500 ml/g/d	50%
Total fish ⁹⁰	Reduce	≥1×/wk vs no intake	2-3 times per week	33%-50%
Food allergy				
Vitamin C ¹¹⁸	Increase	102-201 (mg/d) ²	85 mg	120%-236%
Copper ¹¹⁸	Increase	>2.6 (mg/d) ²	1 mg	260%
Vegetable ¹¹⁵	Reduce	>324 g/d (21/2 cups) vs 153-324 g/d (1-2.5 cups)	3 cups	75% to >100%
Cereal ¹¹⁵	Reduce	Both having less than 143 g/d (5 × 1 oz) and more than 230 g/d (8 × 1 oz)	240 g	<i>Decrease if intake is <60% and >95%</i>
Sensitization				
Vitamin C ¹¹⁸	Increase	130	85 mg	216.7%
Copper ¹¹⁸	Increase	1.6	1 g	160%
Free sugar ¹⁶	Increase	82.4-345.1 g/d	<12 g	687%-2875.8%

Note: Italic = below recommended intake; bold = above recommended intake; normal – stretch across lower than higher than recommended intake.

^aListing only significant OR/RR/HR.

offspring. Four reviews recommended omega-3 fatty acid intake in pregnancy, vitamins A, D, and E; zinc; fruits and vegetables; and a Mediterranean diet for the prevention of asthma.^{43,137-140} Reviews focusing on allergy prevention in general concluded that pregnant women should consume their normal diet and that maternal intake of Mediterranean dietary patterns, diets rich in fruits and vegetables, fish, and vitamin D-containing foods may be associated with reduction of offspring allergic disease.^{10,43,131,132,137,140-146} We were unable to make generalized recommendations from the current review. Another factor that needs to be considered is whether family, paternal, and/or specifically maternal history of allergy play a role in the outcomes measured.

Since the review was performed, one study has shown an association between Diet inflammatory index (DII) scores in pregnancy and childhood asthma outcomes over 10 years.¹⁴⁷ Another study found an association with wheeze trajectories in the child, but not asthma up to 7.5 years of age¹⁴⁸ indicating that perhaps, the inflammatory potential of the diet needs to be assessed when focusing on asthma and/or wheeze outcomes.

4.4 | Interpretation of the evidence

The heterogeneity of the results from observational studies yields little guidance regarding changes in the maternal diet that may be associated with reduction in offspring allergic diseases. Several nutrients, foods and food patterns such as vitamin A, vitamin D, copper, vitamin C, vitamin E, calcium, zinc, beta-carotene, magnesium, dairy/probiotic-containing foods, Mediterranean diet and Western diet,

Mediterranean and soy/fish/nuts diet, fruit and vegetables, fish, nuts, fish/fatty fish, and meat showed associations with reduced allergy outcomes in the observational studies. But amounts consumed do not relate to current nutrition guidelines. Yet, our meta-analysis of RCTs conducted in pregnancy does show some significant and replicable results. Randomized intervention trials in pregnancy show a preventative effect of vitamin D supplementation of 800,⁵⁷ 2400,⁵⁵ or 4000⁶⁰ IU per day on wheeze/asthma; though, EFSA and the IOM recommend only 600 IU of vitamin D per day during pregnancy.

4.5 | Clinical, policy, and research implications of the findings

The current systematic review does not lead to any changes in clinical practice or policies but clearly highlights the research need for more RCTs in this field, particularly focusing on diet patterns or total dietary intake. There may be a particular need to review vitamin D recommendations during pregnancy.

5 | CONCLUSIONS

The lack of consistent results across the different studies presented in this review may largely be influenced by the lack of a standardized approach to supplementation in terms of dose and duration, disease definition/outcomes used, and individual host features that are difficult to compare across studies. Polymorphisms in genes associated

TABLE 3 Summary table for systematic review

Observational trials	Decreased			Increased		
	Vit/Min	Fats/fatty fish	Diet Pattern	Vit/Min	Fats/fatty fish	Diet Pattern
Food allergy	Copper Vitamin C	NA	NA	Vitamin D	NA	Vegetables
Atopic dermatitis	Vitamin E Copper Calcium Zinc Beta-carotene Magnesium	Cholesterol Arachidonic acid	Green vegetables Dairy Fruit (apples) Fish Probiotic milk	Vitamin D Vitamin C	Vegetable oil Fatty fish Margarine Linoleic acid Omega-6 fatty acid Ratio of omega-3/omega-6 Alpha-linolenic acid Total fat Monounsaturated fatty acids	Alcohol Shellfish Meat Fast food Natto
Asthma/wheeze	Vitamin E Zinc Copper Manganese Folic acid Vitamin D Vitamin C Alpha-tocopherol Calcium Vitamin K	DHA Fatty fish Palmitic acid Arachidonic acid Alpha-linolenic acid Saturated fatty acid Olive oil Omega-3 fatty acid	Vegetables Fruit (apples)	Folic acid Vitamin D Vitamin A Vitamin K Alpha-tocopherol Vitamin C	Omega-3 fatty acid Alpha-linolenic acid	Fruit Vegetables
Rhinoconjunctivitis	Vitamin D Vitamin A	Fatty fish Polyunsaturated fatty acid Alpha-linolenic acid	Alcohol Fish Probiotic foods Peanut and pistachio	Vitamin D	Butter Omega-6/omega-3 ratio	Fruit
Sensitization	Vitamin C Copper Vitamin D	Butter Omega-6 fatty acids Polyunsaturated fatty acid	fish	NA	Vegetable fat Oily fish	Fruit Vegetable (celery and sweet pepper) Free sugar
Decreased						
RCTs	Vit/Min	Fats/fatty fish	Foods	Vit/Min	Fats/fatty fish	Foods
Food allergy	NA	No effect	NA	NA	No effect	NA
Atopic dermatitis	No effect	No effect	NA	No effect	No effect	NA
Asthma/wheeze	Vitamin D	No effect	NA	NA	No effect	NA
Rhinoconjunctivitis	NA	No effect	NA	NA	NA	NA
Sensitization	No effect	No effect	NA	No effect	No effect	NA
Increased						
RCTs	Vit/Min	Fats/fatty fish	Foods	Vit/Min	Fats/fatty fish	Foods
Food allergy	NA	No effect	NA	NA	No effect	NA
Atopic dermatitis	No effect	No effect	NA	No effect	No effect	NA
Asthma/wheeze	Vitamin D	No effect	NA	NA	No effect	NA
Rhinoconjunctivitis	NA	No effect	NA	NA	NA	NA
Sensitization	No effect	No effect	NA	No effect	No effect	NA

Note: Italic = only associated with reduced outcomes.

with catabolism and utilization will influence nutrient requirements and function. GWAS-led prevention and intervention studies, including functional microbiome, immunologic, metabolomic, and lipidomic assessments, are required and will increase our understanding of the importance of specific nutrients in the natural course of allergies and asthma. It is likely that a custom-individual-tailored approach to nutrition, focusing on overall dietary intake which may include supplementation, is required to observe the optimal benefits that can potentially be derived from dietary factors in the prevention and treatment of allergies and asthma. Future research and clinical efforts should be focused on large, adequately powered human studies of rigorous methodological design. These studies should focus on identifying the key host characteristics (ie, genetics, environmental factors, microbiome, biochemical and inflammatory parameters, and functional clinical characterization) that influence responses, while also taking levels of nutrient intake, history of allergic diseases, and the composition of the total underlying diet and nutrient interactions into account. Improving our understanding of potential strategies to prevent the development of allergic diseases would significantly reduce morbidity, mortality, and costs from allergic diseases. The current evidence mainly focused on single dietary factors, and the data from observational studies were very heterogeneous. Meta-analysis based on 3 RCTs showed some benefits of vitamin D supplementation on asthma/wheeze, but doses used were higher than recommended by EFSA and the IOM. Before routine supplementation during pregnancy for the prevention of allergic disease can be advised, more studies are required.

CONFLICT OF INTEREST

Venter C provided educational material or reviewed educational materials for Abbott Laboratories, Danone, and Reckitt Benckiser. Agostoni C is an advisor for Ferrero and Dicofarm. Fleischer D is a consultant to Aquestive, Aravax, Genentech, Nasus, AllerGenis, Intrommune, and DOTS Technology Corp. He has provided educational material for Nutricia. He has the following organizational declarations: DBV Technologies—Clinical Medical Advisory Board; Food Allergy & Anaphylaxis Connection Team—Medical Advisory Board; Food Allergy Research & Education—Clinical Advisory Board; and National Peanut Board—Allergy Education Advisory Council. Greenhawt M is supported by grant #5K08HS024599-02 from the Agency for Healthcare Research and Quality; is an expert panel and coordinating committee member of the NIAID-sponsored Guidelines for Peanut Allergy Prevention; has served as a consultant for the Canadian Transportation Agency, Thermo Fisher, Intrommune, and Aimmune Therapeutics; is a member of physician/medical advisory boards for Aimmune Therapeutics, DBV Technologies, Sanofi Genzyme, Genentech, Nutricia, Kaleo Pharmaceutical, Nestle, Aquestive, Allergy Therapeutics, AllerGenis, Aravax, GlaxoSmithKline, Prota, and Monsanto; is a member of the scientific advisory council for the National Peanut Board; has received honorarium for lectures from Thermo Fisher, Aimmune, DBV, Before Brands, multiple state allergy societies, the American College of Allergy, Asthma, and Immunology, and the European Academy of Allergy and Clinical Immunology; is an associate editor for the *Annals of Allergy, Asthma,*

and Immunology; and is a member of the Joint Task Force on Allergy Practice Parameters. Meyer R has grants from Danone, personal fees from Nutricia and Mead Johnson, and is an advisor to CoMISS group from Nestle. Netting MJ has received honorarium to her institution for speaker fees from Nestle. Smith PK has provided lecture material and/or consultancy to Abbot, NNI, and Danone. He is a PI for DBV and has received investigator-initiated funding from Nestle. O'Mahony L has provided consultancy with Alimentary Health and research grant from GSK. The following authors have no conflict of interest: Vanderlinden L, Untersmayr E, Roberts G, Roduit C, Palumbo MP, Palmer DJ, Nwaru BI, Muraro A, Maiorella A, Maslin K, Lunjani N, Groetch M, Glueck DH, Du Toit G, Ben-Abdallah M, and Arshad SH

AUTHOR CONTRIBUTION

CV drafted the paper. MBA, MP, DG, AMai, LV people performed the data extraction and analysis. CA, DF, MG, EU, GR, CR, DP, AMu, KM, NL, MG, GdT, SHA, PKS, MN, RM reviewed and contributed to multiple versions of the paper. LOM and BN assisted with data interpretation and reviewed and edited the final version of the paper.

ORCID

Carina Venter  <https://orcid.org/0000-0002-7473-5355>
 Carlo Agostoni  <https://orcid.org/0000-0002-5006-0832>
 S. Hasan Arshad  <https://orcid.org/0000-0001-5988-235X>
 George Du Toit  <https://orcid.org/0000-0002-0321-2928>
 Matthew Greenhawt  <https://orcid.org/0000-0002-2365-9372>
 Deborah H. Glueck  <https://orcid.org/0000-0003-0047-8939>
 Marion Groetch  <https://orcid.org/0000-0002-5931-881X>
 Kate Maslin  <https://orcid.org/0000-0001-5580-803X>
 Rosan Meyer  <https://orcid.org/0000-0002-5710-5570>
 Muraro Antonella  <https://orcid.org/0000-0002-5026-5862>
 Merryn J. Netting  <https://orcid.org/0000-0002-0642-9145>
 Bright Ibeabughichi Nwaru  <https://orcid.org/0000-0002-2876-6089>
 Debra J. Palmer  <https://orcid.org/0000-0003-2498-8343>
 Micheala P. Palumbo  <https://orcid.org/0000-0002-0267-5779>
 Graham Roberts  <https://orcid.org/0000-0003-2252-1248>
 Caroline Roduit  <https://orcid.org/0000-0002-5988-0570>
 Pete Smith  <https://orcid.org/0000-0002-2629-8963>
 Eva Untersmayr  <https://orcid.org/0000-0002-1963-499X>
 Lauren A. Vanderlinden  <https://orcid.org/0000-0002-4019-8395>
 Liam O'Mahony  <https://orcid.org/0000-0003-4705-3583>
 Deborah H. Glueck  <https://orcid.org/0000-0003-0047-8939>

REFERENCES

1. Institute of Medicine. Food Allergies: Global Burden, Causes, Treatment, Prevention and Public Policy. National Academy of Sciences. <http://www.nationalacademies.org/hmd/Activities/Nutrition/FoodAllergies.aspx>. Published 2016. Updated 28 March 2017. Accessed March, 2017.
2. World Allergy Organization. World Allergy Organization White Book on Allergy. 2011: https://www.worldallergy.org/UserFiles/file/WAO-White-Book-on-Allergy_web.pdf

3. Pawankar RCG, Holgate ST, Lockey RF, Blaiss M. *The WAO White Book on Allergy*. 2013. <https://www.worldallergy.org/UserFiles/file/WhiteBook2-2013-v8.pdf>
4. Agache I, Akdis CA. Precision medicine and phenotypes, endotypes, genotypes, regiotypes, and theratypes of allergic diseases. *J Clin Invest*. 2019;130:1493-1503.
5. Muraro A, Dubois AEJ, DunnGalvin A, et al. EAACI Food Allergy and Anaphylaxis Guidelines. Food allergy health-related quality of life measures. *Allergy*. 2014;69(7):845-853.
6. Dwyer-Lindgren L, Bertozzi-Villa A, Stubbs RW, et al. Trends and patterns of differences in chronic respiratory disease mortality among US counties, 1980-2014. *JAMA*. 2017;318(12):1136-1149.
7. Turner PJ, Gowland MH, Sharma V, et al. Increase in anaphylaxis-related hospitalizations but no increase in fatalities: an analysis of United Kingdom national anaphylaxis data, 1992-2012. *J Allergy Clin Immunol*. 2015;135(4):956-963.e1.
8. Venter C, Greenhawt M, Meyer RW, et al. EAACI position paper on diet diversity in pregnancy, infancy and childhood: Novel concepts and implications for studies in allergy and asthma. *Allergy*. 2019;75(3):497-523.
9. Venter C, Meyer RW, Nwaru BI, et al. EAACI position paper: Influence of dietary fatty acids on asthma, food allergy, and atopic dermatitis. *Allergy*. 2019;74(8):1429-1444.
10. de Silva D, Geromi M, Halken S, et al. Primary prevention of food allergy in children and adults: systematic review. *Allergy*. 2014;69(5):581-589.
11. de Silva D, Halken S, Singh C, et al. Preventing immediate-onset food allergy in infants, children and adults: Systematic review protocol. *Pediatr Allergy Immunol*. 2020;31(3):243-249.
12. Venter C, Brown KR, Maslin K, Palmer DJ. Maternal dietary intake in pregnancy and lactation and allergic disease outcomes in offspring. *Pediatr Allergy Immunol*. 2017;28(2):135-143.
13. Garcia-Larsen V, Ierodiakonou D, Jarrold K, et al. Diet during pregnancy and infancy and risk of allergic or autoimmune disease: A systematic review and meta-analysis. *PLoS Medicine*. 2018;15(2):e1002507.
14. Alfonso VH, Bandoli G, von Ehrenstein O, Ritz B. Early folic acid Supplement initiation and risk of adverse early childhood respiratory health: a population-based study. *Matern Child Health J*. 2018;22(1):111-119.
15. Allan KM, Prabhu N, Craig LC, et al. Maternal vitamin D and E intakes during pregnancy are associated with asthma in children. *Eur Respir J*. 2015;45(4):1027-1036.
16. Bedard A, Northstone K, Henderson AJ, Shaheen SO. Maternal intake of sugar during pregnancy and childhood respiratory and atopic outcomes. *Eur Respir J*. 2017;50(1).
17. Bedard A, Northstone K, Holloway JW, Henderson AJ, Shaheen SO. Maternal dietary antioxidant intake in pregnancy and childhood respiratory and atopic outcomes: birth cohort study. *Eur Respir J*. 2018;52(2):1800507.
18. Bunyavanich S, Rifas-Shiman SL, Platts-Mills TA, et al. and childhood vitamin D exposure and their association with childhood allergic rhinitis and allergic sensitization. *J Allergy Clin Immunol*. 2016;137(4):1063-1070.e2.
19. Castro-Rodriguez JA, Ramirez-Hernandez M, Padilla O, Pacheco-Gonzalez RM, Perez-Fernandez V, Garcia-Marcos L. Effect of foods and Mediterranean diet during pregnancy and first years of life on wheezing, rhinitis and dermatitis in preschoolers. *Allergol Immunopathol*. 2016;44(5):400-409.
20. Chatzi L, Garcia R, Roumeliotaki T, et al. Mediterranean diet adherence during pregnancy and risk of wheeze and eczema in the first year of life: INMA (Spain) and RHEA (Greece) mother-child cohort studies. *Br J Nutr*. 2013;110(11):2058-2068.
21. de Batlle J, Garcia-Aymerich J, Barraza-Villarreal A, Anto JM, Romieu I. Mediterranean diet is associated with reduced asthma and rhinitis in Mexican children. *Allergy*. 2008;63(10):1310-1316.
22. den Dekker HT, Jaddoe VVW, Reiss IK, de Jongste JC, Duijts L. Maternal folic acid use during pregnancy, methylenetetrahydrofolate reductase gene polymorphism, and child's lung function and asthma. *Clin Exp Allergy*. 2018;48(2):175-185.
23. Devereux G, Craig L, Seaton A, Turner S. Maternal vitamin D and E intakes in pregnancy and asthma to age 15 years: a cohort study. *Pediatr Pulmonol*. 2019;54(1):11-19.
24. Loo EXL, Ong L, Goh A, et al. Effect of Maternal Dietary Patterns during Pregnancy on Self-Reported Allergic Diseases in the First 3 Years of Life: Results from the GUSTO Study. *Int Arch Allergy Immunol*. 2017;173(2):105-113.
25. Maslova E, Hansen S, Strom M, Halldorsson TI, Olsen SF. Maternal intake of vitamins A, E and K in pregnancy and child allergic disease: a longitudinal study from the Danish National Birth Cohort. *Br J Nutr*. 2014;111(6):1096-1108.
26. Miyake Y, Tanaka K, Okubo H, Sasaki S, Arakawa M. Maternal consumption of dairy products, calcium, and vitamin D during pregnancy and infantile allergic disorders. *Ann Allergy Asthma Immunol*. 2014;113(1):82-87.
27. Moonesinghe H, Patil VK, Dean T, et al. Association between healthy eating in pregnancy and allergic status of the offspring in childhood. *Ann Allergy Asthma Immunol*. 2016;116(2):163-165.
28. Ogawa K, Morisaki N, Kobayashi M, et al. Maternal vegetable intake in early pregnancy and wheeze in offspring at the age of 2 years. *Eur J Clin Nutr*. 2018;72(5):761-771.
29. Ozawa N, Shimojo N, Suzuki Y, et al. Maternal intake of Natto, a Japan's traditional fermented soybean food, during pregnancy and the risk of eczema in Japanese babies. *Allergol Int*. 2014;63(2):261-266.
30. Parr CL, Magnus MC, Karlstad O, et al. Maternal folate intake during pregnancy and childhood asthma in a population-based cohort. *Am J Respir Crit Care Med*. 2017;195(2):221-228.
31. Shaheen SO, Rutterford C, Zuccolo L, et al. Prenatal alcohol exposure and childhood atopic disease: a Mendelian randomization approach. *J Allergy Clin Immunol*. 2014;133(1):225-232.e5.
32. Stratakis N, Roumeliotaki T, Oken E, et al. Fish and seafood consumption during pregnancy and the risk of asthma and allergic rhinitis in childhood: a pooled analysis of 18 European and US birth cohorts. *Int J Epidemiol*. 2017;46(5):1465-1477.
33. Trivedi MK, Sharma S, Rifas-Shiman SL, et al. Folic Acid in Pregnancy and Childhood Asthma: A US Cohort. *Clin Pediatr (Phila)*. 2018;57(4):421-427.
34. Tuokkola J, Luukkainen P, Kaila M, et al. Maternal dietary folate, folic acid and vitamin D intakes during pregnancy and lactation and the risk of cows' milk allergy in the offspring. *Br J Nutr*. 2016;116(4):710-718.
35. Veeranki SP, Gebretsadik T, Mitchel EF, et al. Maternal folic acid supplementation during pregnancy and early childhood asthma. *Epidemiology*. 2015;26(6):934-941.
36. Viljoen K, Segurado R, O'Brien J, et al. Pregnancy diet and offspring asthma risk over a 10-year period: the lifeways cross generation cohort study, Ireland. *BMJ Open*. 2018;8(2):e017013.
37. Wada K, Konishi K, Tamura T, Shiraki M, Iwasa S, Nagata C. Alcohol intake during pregnancy and offspring's atopic eczema risk. *Alcohol Clin Exp Res*. 2016;40(5):1037-1043.
38. Yang L, Jiang L, Bi M, et al. High dose of maternal folic acid supplementation is associated to infant asthma. *Food Chem Toxicol*. 2015;75:88-93.
39. Best KP, Sullivan TR, Palmer DJ, et al. Prenatal omega-3 LCPUFA and symptoms of allergic disease and sensitization throughout early childhood - a longitudinal analysis of long-term follow-up of a randomized controlled trial. *World Allergy Organ J*. 2018;11(1):10.
40. Wolsk HM, Chawes BL, Litonjua AA, et al. Prenatal vitamin D supplementation reduces risk of asthma/recurrent wheeze in early childhood: A combined analysis of two randomized controlled trials. *PLoS One*. 2017;12(10):e0186657.

41. Allergy ASocla. Australasian Society of Clinical Immunology and Allergy guidelines for infant feeding and allergy prevention. <https://www.allergy.org.au/patients/allergy-prevention/ascia-guidelines-for-infant-feeding-and-allergy-prevention> Published 2016. Accessed.
42. British Dietetic Association. Early Feeding Guidance. <https://www.bsaci.org/about/early-feeding-guidance>. Published 2018. Accessed April, 2019.
43. Muraro A, Halken S, Arshad SH, et al. EAACI Food Allergy and Anaphylaxis Guidelines. Primary prevention of food allergy. *Allergy*. 2014;69(5):590-601.
44. Greer FR, Sicherer SH, Burks AW. The effects of early nutritional interventions on the development of atopic disease in infants and children: the role of maternal dietary restriction, breastfeeding, hydrolyzed formulas, and timing of introduction of allergenic complementary foods. *Pediatrics*. 2019;143(4):e20190281.
45. Schafer T, Bauer CP, Beyer K, et al. S3-Guideline on allergy prevention: 2014 update: Guideline of the German Society for Allergology and Clinical Immunology (DGAKI) and the German Society for Pediatric and Adolescent Medicine (DGKJ). *Allergo J Int*. 2014;23(6):186-199.
46. Maslova E, Granstrom C, Hansen S, et al. Peanut and tree nut consumption during pregnancy and allergic disease in children-should mothers decrease their intake? Longitudinal evidence from the Danish National Birth Cohort. *J Allergy Clin Immunol*. 2012;130(3):724-732.
47. du Toit G, Sayre PH, Roberts G, et al. The allergen-specificity of early peanut consumption and the impact on the development of allergic disease in the LEAP Study Cohort. *J Allergy Clin Immunol*. 2018;141(4):1343-1353.
48. National Academies Press. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids. 2002.
49. National Institute of Health and Care Excellence. Methods for development of NICE public health guidance. <https://www.nice.org.uk/process/pmg4/chapter/introduction>. Accessed April, 2019.
50. Grade Handbook. Introduction to Grade Handbook. <https://gdt.gradepro.org/app/handbook/handbook.html#h.ygojbnr1bi5y>. Published 2019. Accessed August 2019.
51. T. L. Meta-Analysis. R package version 3.0. <https://CRAN.R-project.org/package=rmeta>. Published 2018. Accessed August 2019.
52. Gordon M, L. T. Advanced Forest Plot Using 'grid' Graphics. R package version 1.9. <https://CRAN.R-project.org/package=forestplot>. Published 2019. Accessed 2019.
53. Berman D, Clinton C, Limb R, Somers EC, Romero V, Mozurkewich E. Prenatal omega-3 supplementation and eczema risk among offspring at age 36 months. *Insights Allergy Asthma Bronchitis*. 2016;2(1). <https://doi.org/10.21767/2471-304X.100014>
54. Bisgaard H, Stokholm J, Chawes BL, et al. Fish oil-derived fatty acids in pregnancy and wheeze and asthma in offspring. *N Engl J Med*. 2016;375(26):2530-2539.
55. Chawes BL, Bonnelykke K, Stokholm J, et al. Effect of vitamin D3 supplementation during pregnancy on risk of persistent wheeze in the offspring: a randomized clinical trial. *JAMA*. 2016;315(4):353-361.
56. Dunstan JA, Mori TA, Barden A, et al. Fish oil supplementation in pregnancy modifies neonatal allergen-specific immune responses and clinical outcomes in infants at high risk of atopy: a randomized, controlled trial. *J Allergy Clin Immunol*. 2003;112(6):1178-1184.
57. Goldring ST, Griffiths CJ, Martineau AR, et al. Prenatal vitamin D supplementation and child respiratory health: a randomised controlled trial. *PLoS One*. 2013;8(6):e66627.
58. Greenough A, Shaheen SO, Shennan A, Seed PT, Poston L. Respiratory outcomes in early childhood following antenatal vitamin C and E supplementation. *Thorax*. 2010;65(11):998-1003.
59. Imhoff-Kunsch B, Stein AD, Martorell R, Parra-Cabrera S, Romieu I, Ramakrishnan U. Prenatal docosahexaenoic acid supplementation and infant morbidity: randomized controlled trial. *Pediatrics*. 2011;128(3):505-512.
60. Litonjua AA, Carey VJ, Laranjo N, et al. Effect of prenatal supplementation with vitamin D on asthma or recurrent wheezing in offspring by age 3 years: the VDAART randomized clinical trial. *JAMA*. 2016;315(4):362-370.
61. McEvoy CT, Schilling D, Clay N, et al. Vitamin C supplementation for pregnant smoking women and pulmonary function in their newborn infants: a randomized clinical trial. *JAMA*. 2014;311(20):2074-2082.
62. Noakes PS, Vlachava M, Kremmyda LS, Diaper ND, Miles EA, Erlewyn-Lajeunesse M. Increased intake of oily fish in pregnancy: Effects on neonatal immune responses and on clinical outcomes in infants at 6 mo. *Am J Clin Nutr*. 2012;95(2):395-404.
63. Olsen SF, Osterdal ML, Salvig JD, et al. Fish oil intake compared with olive oil intake in late pregnancy and asthma in the offspring: 16 y of registry-based follow-up from a randomized controlled trial. *Am J Clin Nutr*. 2008;88(1):167-175.
64. Hansen S, Strom M, Maslova E, et al. Fish oil supplementation during pregnancy and allergic respiratory disease in the adult offspring. *J Allergy Clin Immunol*. 2017;139(1):104-111.e4.
65. Palmer DJ, Sullivan T, Gold MS, et al. Effect of n-3 long chain polyunsaturated fatty acid supplementation in pregnancy on infants' allergies in first year of life: randomised controlled trial. *BMJ*. 2012;344:184.
66. Palmer DJ, Sullivan T, Gold MS, et al. Randomized controlled trial of fish oil supplementation in pregnancy on childhood allergies. *Allergy*. 2013;68(11):1370-1376.
67. Best KP, Sullivan T, Palmer D, et al. Prenatal fish oil supplementation and allergy: 6-year follow-up of a randomized controlled trial. *Pediatrics*. 2016;137(6):e20154443.
68. Bekkers MB, Elstgeest LE, Scholtens S, et al. Maternal use of folic acid supplements during pregnancy, and childhood respiratory health and atopy. *Eur Respir J*. 2012;39(6):1468-1474.
69. Willers SM, Wijga AH, Brunekreef B, et al. Maternal food consumption during pregnancy and the longitudinal development of childhood asthma. *Am J Respir Crit Care Med*. 2008;178(2):124-131.
70. Bertelsen RJ, Brantsaeter AL, Magnus MC, et al. Probiotic milk consumption in pregnancy and infancy and subsequent childhood allergic diseases. *J Allergy Clin Immunol*. 2014;133(1):165-171.e8.
71. Binkley KE, Leaver C, Ray JG. Antenatal risk factors for peanut allergy in children. *Allergy Asthma Clin Immunol*. 2011;7(1):17.
72. Bisgaard H, Halkjaer LB, Hinge R, et al. Risk analysis of early childhood eczema. *J Allergy Clin Immunol*. 2009;123(6):1355-1360.e1355.
73. Calvani M, Alessandri C, Sopo SM, et al. Consumption of fish, butter and margarine during pregnancy and development of allergic sensitizations in the offspring: role of maternal atopy. *Pediatr Allergy Immunol*. 2006;17(2):94-102.
74. Castro-Rodriguez JA, Garcia-Marcos L, Sanchez-Solis M, Perez-Fernandez V, Martinez-Torres A, Mallol J. Olive oil during pregnancy is associated with reduced wheezing during the first year of life of the offspring. *Pediatr Pulmonol*. 2010;45(4):395-402.
75. Chatzi L, Torrent M, Romieu I, et al. Mediterranean diet in pregnancy is protective for wheeze and atopy in childhood. *Thorax*. 2008;63(6):507-513.
76. Romieu I, Torrent M, Garcia-Esteban R, et al. Maternal fish intake during pregnancy and atopy and asthma in infancy. *Clin Exp Allergy*. 2007;37(4):518-525.
77. Dunstan JA, West C, McCarthy S, et al. The relationship between maternal folate status in pregnancy, cord blood

- folate levels, and allergic outcomes in early childhood. *Allergy*. 2012;67(1):50-57.
78. Fitzsimon N, Fallon U, O'Mahony D, et al. Mothers' dietary patterns during pregnancy and risk of asthma symptoms in children at 3 years. *Ir Med J*. 2007;100(8):27-32.
 79. Haberg SE, London SJ, Stigum H, Nafstad P, Nystad W. Folic acid supplements in pregnancy and early childhood respiratory health. *Arch Dis Child*. 2009;94(3):180-184.
 80. Jedrychowski W, Flak E, Mroz E, et al. Modulating effects of maternal fish consumption on the occurrence of respiratory symptoms in early infancy attributed to prenatal exposure to fine particles. *Ann Nutr Metab*. 2008;52(1):8-16.
 81. Jedrychowski W, Perera F, Maugeri U, et al. Effects of prenatal and perinatal exposure to fine air pollutants and maternal fish consumption on the occurrence of infantile eczema. *Int Arch Allergy Immunol*. 2011;155(3):275-281.
 82. Lange NE, Rifas-Shiman SL, Camargo Jr CA, Gold DR, Gillman MW, Litonjua AA. Maternal dietary pattern during pregnancy is not associated with recurrent wheeze in children. *J Allergy Clin Immunol*. 2010;126(2):250-255, 255 e251-254.
 83. Litonjua AA, Rifas-Shiman SL, Ly NP, et al. Maternal antioxidant intake in pregnancy and wheezing illnesses in children at 2 y of age. *Am J Clin Nutr*. 2006;84(4):903-911.
 84. Camargo CA Jr, Rifas-Shiman SL, Litonjua AA, et al. Maternal intake of vitamin D during pregnancy and risk of recurrent wheeze in children at 3 y of age. *Am J Clin Nutr*. 2007;85(3):788-795.
 85. Leermakers ET, Sonnenschein-van der Voort AM, Heppe DH, et al. Maternal fish consumption during pregnancy and risks of wheezing and eczema in childhood: the Generation R Study. *Eur J Clin Nutr*. 2013;67(4):353-359.
 86. Magdelijns FJ, Mommers M, Penders J, Smits L, Thijs C. Folic acid use in pregnancy and the development of atopy, asthma, and lung function in childhood. *Pediatrics*. 2011;128(1):e135-e144.
 87. Martindale S, McNeill G, Devereux G, Campbell D, Russell G, Seaton A. Antioxidant intake in pregnancy in relation to wheeze and eczema in the first two years of life. *Am J Respir Crit Care Med*. 2005;171(2):121-128.
 88. Devereux G, Turner SW, Craig LC, et al. Low maternal vitamin E intake during pregnancy is associated with asthma in 5-year-old children. *Am J Respir Crit Care Med*. 2006;174(5):499-507.
 89. Devereux G, Litonjua AA, Turner SW, et al. Maternal vitamin D intake during pregnancy and early childhood wheezing. *Am J Clin Nutr*. 2007;85(3):853-859.
 90. Willers SM, Devereux G, Craig LC, et al. Maternal food consumption during pregnancy and asthma, respiratory and atopic symptoms in 5-year-old children. *Thorax*. 2007;62(9):773-779.
 91. Maslova E, Hansen S, Jensen CB, Thorne-Lyman AL, Strom M, Olsen SF. Vitamin D intake in mid-pregnancy and child allergic disease - a prospective study in 44,825 Danish mother-child pairs. *BMC Pregnancy Childbirth*. 2013;13:199.
 92. Maslova E, Strom M, Olsen SF, Halldorsson TI. Consumption of artificially-sweetened soft drinks in pregnancy and risk of child asthma and allergic rhinitis. *PLoS One*. 2013;8(2):e57261.
 93. Maslova E, Strom M, Oken E, et al. Fish intake during pregnancy and the risk of child asthma and allergic rhinitis - longitudinal evidence from the Danish National Birth Cohort. *Br J Nutr*. 2013;110(7):1313-1325.
 94. Linneberg A, Petersen J, Gronbaek M, Benn CS. Alcohol during pregnancy and atopic dermatitis in the offspring. *Clin Exp Allergy*. 2004;34(11):1678-1683.
 95. Miyake Y, Tanaka K, Okubo H, Sasaki S, Arakawa M. Maternal fat intake during pregnancy and wheeze and eczema in Japanese infants: the Kyushu Okinawa Maternal and Child Health Study. *Ann Epidemiol*. 2013;23(11):674-680.
 96. Miyake Y, Sasaki S, Tanaka K, Ohfuji S, Hirota Y. Maternal fat consumption during pregnancy and risk of wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. *Thorax*. 2009;64(9):815-821.
 97. Saito K, Yokoyama T, Miyake Y, et al. Maternal meat and fat consumption during pregnancy and suspected atopic eczema in Japanese infants aged 3-4 months: the Osaka Maternal and Child Health Study. *Pediatr Allergy Immunol*. 2010;21(1 Pt 1):38-46.
 98. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Dairy food, calcium and vitamin D intake in pregnancy, and wheeze and eczema in infants. *Eur Respir J*. 2010;35(6):1228-1234.
 99. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Consumption of vegetables, fruit, and antioxidants during pregnancy and wheeze and eczema in infants. *Allergy*. 2010;65(6):758-765.
 100. Miyake Y, Sasaki S, Tanaka K, Hirota Y. Maternal B vitamin intake during pregnancy and wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. *Pediatr Allergy Immunol*. 2011;22(1 Pt 1):69-74.
 101. Miyake Y, Okubo H, Sasaki S, Tanaka K, Hirota Y. Maternal dietary patterns during pregnancy and risk of wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. *Pediatr Allergy Immunol*. 2011;22(7):734-741.
 102. Nwaru BI, Ahonen S, Kaila M, et al. Maternal diet during pregnancy and allergic sensitization in the offspring by 5 yrs of age: a prospective cohort study. *Pediatr Allergy Immunol*. 2010;21(1 Pt 1):29-37.
 103. Nwaru BI, Erkkola M, Ahonen S, et al. Intake of antioxidants during pregnancy and the risk of allergies and asthma in the offspring. *Eur J Clin Nutr*. 2011;65(8):937-943.
 104. Nwaru BI, Erkkola M, Lumia M, et al. Maternal intake of fatty acids during pregnancy and allergies in the offspring. *Br J Nutr*. 2012;108(4):720-732.
 105. Lumia M, Luukkainen P, Tapanainen H, et al. Dietary fatty acid composition during pregnancy and the risk of asthma in the offspring. *Pediatr Allergy Immunol*. 2011;22(8):827-835.
 106. Erkkola M, Nwaru BI, Kaila M, et al. Risk of asthma and allergic outcomes in the offspring in relation to maternal food consumption during pregnancy: a Finnish birth cohort study. *Pediatr Allergy Immunol*. 2012;23(2):186-194.
 107. Erkkola M, Kaila M, Nwaru BI, et al. Maternal vitamin D intake during pregnancy is inversely associated with asthma and allergic rhinitis in 5-year-old children. *Clin Exp Allergy*. 2009;39(6):875-882.
 108. Oien T, Storro O, Johnsen R. Do early intake of fish and fish oil protect against eczema and doctor-diagnosed asthma at 2 years of age? A cohort study. *J Epidemiol Community Health*. 2010;64(2):124-129.
 109. Oliveti JF, Kerckmar CM, Redline S. Pre- and perinatal risk factors for asthma in inner city African-American children. *Am J Epidemiol*. 1996;143(6):570-577.
 110. Pike KC, Inskip HM, Robinson S, et al. Maternal late-pregnancy serum 25-hydroxyvitamin D in relation to childhood wheeze and atopic outcomes. *Thorax*. 2012;67(11):950-956.
 111. Salam MT, Li YF, Langholz B, Gilliland FD. Maternal fish consumption during pregnancy and risk of early childhood asthma. *J Asthma*. 2005;42(6):513-518.
 112. Sausenthaler S, Koletzko S, Schaaf B, et al. Maternal diet during pregnancy in relation to eczema and allergic sensitization in the offspring at 2 y of age. *Am J Clin Nutr*. 2007;85(2):530-537.
 113. Shaheen SO, Northstone K, Newson RB, Emmett PM, Sherriff A, Henderson AJ. Dietary patterns in pregnancy and respiratory and atopic outcomes in childhood. *Thorax*. 2009;64(5):411-417.
 114. Granell R, Heron J, Lewis S, Davey Smith G, Sterne JA, Henderson J. The association between mother and child MTHFR C677T polymorphisms, dietary folate intake and childhood atopy in a population-based, longitudinal birth cohort. [Erratum appears in Clin Exp

- Allergy. 2008 Apr; 38(4):699 Note: Smith, G D [corrected to Davey Smith, G]]. *Clin Exp Allergy*. 2008;38(2):320-328.
115. Tuokkola J, Luukkainen P, Tapanainen H, et al. Maternal diet during pregnancy and lactation and cow's milk allergy in offspring. *Eur J Clin Nutr*. 2016;70(5):554-559.
 116. Wang J, Guo YL, Weng HJ, et al. Environmental risk factors for early infantile atopic dermatitis. *Pediatr Allergy Immunol*. 2007;18(5):441-447.
 117. Watson PE, McDonald BW. Water and nutrient intake in pregnant New Zealand women: association with wheeze in their infants at 18 months. *Asia Pacific J Clin Nutr*. 2014;23(4):660-670.
 118. West CE, Dunstan J, McCarthy S, et al. Associations between maternal antioxidant intakes in pregnancy and infant allergic outcomes. *Nutrients*. 2012;4(11):1747-1758.
 119. Whitrow MJ, Moore VM, Rumbold AR, Davies MJ. Effect of supplemental folic acid in pregnancy on childhood asthma: a prospective birth cohort study. *Am J Epidemiol*. 2009;170(12):1486-1493.
 120. Ioannidis JP. Implausible results in human nutrition research. *BMJ*. 2013;347:f6698.
 121. Scott M, Kurukulaarachy RJ, Raza A, Arshad SH. Understanding the nature and outcome of childhood wheezing. *Eur Respir J*. 2009;33(3):700-701.
 122. Raza A, Kurukulaarachy RJ, Grundy JD, et al. What does adolescent undiagnosed wheeze represent? Findings from the Isle of Wight Cohort. *Eur Respir J*. 2012;40(3):580-588.
 123. European Academy of Allergy Clinical Immunology. Global atlas of Allergy. <http://www.eaaci.org/globalatlas/GlobalAtlasAllergy.pdf>. Published 2014. Accessed 2018.
 124. Kominarek MA, Rajan P. Nutrition Recommendations in Pregnancy and Lactation. *Med Clin North Am*. 2016;100(6):1199-1215.
 125. Norizoe C, Akiyama N, Segawa T, et al. Increased food allergy and vitamin D: randomized, double-blind, placebo-controlled trial. *Pediatr Int*. 2014;56(1):6-12.
 126. Vahdaninia M, Mackenzie H, Dean T, Helps S. omega-3 LCPUFA supplementation during pregnancy and risk of allergic outcomes or sensitization in offspring: A systematic review and meta-analysis. *Ann Allergy Asthma Immunol*. 2019;122(3):302-313 e302.
 127. Furuholm C, Warstedt K, Fageras M, et al. Allergic disease in infants up to 2 years of age in relation to plasma omega-3 fatty acids and maternal fish oil supplementation in pregnancy and lactation. *Pediatr Allergy Immunol*. 2011;22(5):505-514.
 128. Azad MB, Coneys JG, Kozyrskyj AL, et al. Probiotic supplementation during pregnancy or infancy for the prevention of asthma and wheeze: systematic review and meta-analysis. *BMJ*. 2013;347:f6471.
 129. Cuello-Garcia CA, Brozek JL, Fiocchi A, et al. Probiotics for the prevention of allergy: a systematic review and meta-analysis of randomized controlled trials. *J Allergy Clin Immunol*. 2015;136(4):952-961.
 130. Doege K, Grajecki D, Zyriax BC, Detinkina E, Zu Eulenburg C, Buhling KJ. Impact of maternal supplementation with probiotics during pregnancy on atopic eczema in childhood—a meta-analysis. *Br J Nutr*. 2012;107(1):1-6.
 131. Lis-Swiety A, Milewska-Wrobel D, Janicka I. Dietary strategies for primary prevention of atopic diseases - what do we know? *Med Wieku Rozwoj*. 2016;20(1):68-74.
 132. Mansfield JA, Bergin SW, Cooper JR, Olsen CH. Comparative probiotic strain efficacy in the prevention of eczema in infants and children: a systematic review and meta-analysis. *Mil Med*. 2014;179(6):580-592.
 133. Pelucchi C, Chatenoud L, Turati F, et al. Probiotics supplementation during pregnancy or infancy for the prevention of atopic dermatitis: a meta-analysis. *Epidemiology*. 2012;23(3):402-414.
 134. Tang ML, Lahtinen SJ, Boyle RJ. Probiotics and prebiotics: clinical effects in allergic disease. *Curr Opin Pediatrics*. 2010;22(5):626-634.
 135. Zuccotti G, Meneghin F, Aceti A, et al. Probiotics for prevention of atopic diseases in infants: systematic review and meta-analysis. *Allergy*. 2015;70(11):1356-1371.
 136. Vahdaninia M, Mackenzie H, Helps S, Dean T. Prenatal intake of vitamins and allergic outcomes in the offspring: a systematic review and meta-analysis. *J Allergy Clin Immunol Pract*. 2017;5(3):771-778.e5.
 137. Lodge CJ, Allen KJ, Lowe AJ, Dharmage SC. Overview of evidence in prevention and aetiology of food allergy: a review of systematic reviews. *Int J Environ Res Public Health*. 2013;10(11):5781-5806.
 138. Lv N, Xiao L, Ma J. Dietary pattern and asthma: a systematic review and meta-analysis. *J Asthma Allergy*. 2014;7:105-121.
 139. Yang H, Xun P, He K. Fish and fish oil intake in relation to risk of asthma: a systematic review and meta-analysis. *PLoS One*. 2013;8(11):e80048.
 140. Nurmatov U, Devereux G, Sheikh A. Nutrients and foods for the primary prevention of asthma and allergy: systematic review and meta-analysis. *J Allergy Clin Immunol*. 2011;127(3):724-733.e30.
 141. Beckhaus AA, Garcia-Marcos L, Forno E, Pacheco-Gonzalez RM, Celedon JC, Castro-Rodriguez JA. Maternal nutrition during pregnancy and risk of asthma, wheeze, and atopic diseases during childhood: a systematic review and meta-analysis. *Allergy*. 2015;70(12):1588-1604.
 142. Best KP, Gold M, Kennedy D, Martin J, Makrides M. Omega-3 long-chain PUFA intake during pregnancy and allergic disease outcomes in the offspring: a systematic review and meta-analysis of observational studies and randomized controlled trials. *Am J Clin Nutr*. 2016;103(1):128-143.
 143. Brown SB, Reeves KW, Bertone-Johnson ER. Maternal folate exposure in pregnancy and childhood asthma and allergy: a systematic review. *Nutr Rev*. 2014;72(1):55-64.
 144. Crider KS, Cordero AM, Qi YP, Mulinare J, Dowling NF, Berry RJ. Prenatal folic acid and risk of asthma in children: a systematic review and meta-analysis. *Am J Clin Nutr*. 2013;98(5):1272-1281.
 145. Gunaratne AW, Makrides M, Collins CT. Maternal prenatal and/or postnatal n-3 long chain polyunsaturated fatty acids (LCPUFA) supplementation for preventing allergies in early childhood. *Cochrane Database Syst Rev*. 2015;7:CD010085.
 146. Netting MJ, Middleton PF, Makrides M. Does maternal diet during pregnancy and lactation affect outcomes in offspring? A systematic review of food-based approaches. *Nutrition*. 2014;30(11-12):1225-1241.
 147. Chen LW, Lyons B, Navarro P, et al. Maternal dietary inflammatory potential and quality are associated with offspring asthma risk over 10-year follow-up: the Lifeways Cross-Generation Cohort Study. *Am J Clin Nutr*. 2020;111(2):440-447.
 148. Hanson C, Rifas-Shiman SL, Shivappa N, et al. Associations of prenatal dietary inflammatory potential with childhood respiratory outcomes in project viva. *J Allergy Clin Immunol Pract*. 2020;8(3):945-945.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Venter C, Agostoni C, Arshad SH, et al. Dietary factors during pregnancy and atopic outcomes in childhood: A systematic review from the European Academy of Allergy and Clinical Immunology. *Pediatr Allergy Immunol*. 2020;31:889-912. <https://doi.org/10.1111/pai.13303>